

IL-4R/CD124(Tyr497) polyclonal antibody

Catalog: AP0210

Host: Rabbit

Reactivity: Human

Background:

The IL-2 receptor is a multicomponent complex consisting of three subunits, α , β and γ , each of which is required for high affinity binding of IL-2. The α chain functions primarily in binding IL-2, whereas the β and γ chains contribute to IL-2 binding and are essential to IL-2-induced activation of signaling pathways leading to T cell growth. Both IL-4R and IL-7R were initially described as single chain, high-affinity ligand-binding cytokine receptors. However, it is now well established that the IL-2R γ chain functions as a second subunit of the high affinity IL-4R and IL-7R receptors. Consequently, the originally described subunits of these latter receptors are now referred to as IL-4R α and IL-7R α , respectively, while the common subunit is referred to as γ c. Although the common γ chain enhances ligand binding in these three cytokine receptors, it has no capacity to bind these ligands on its own. There is evidence that the γ c chain is also a subunit of IL-13R.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.4.

Molecular Weight:

~ 85 kDa

Swiss-Prot:

P24394

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000~1:2000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

IL-4R/CD124(Tyr497) polyclonal antibody detects endogenous levels of HIST1H3A protein.

DATA:



Western blot (WB) analysis of IL-4R/CD124(Tyr497) polyclonal antibody at 1:2000 dilution

Lane1:HepG2 whole cell lysate(40ug)

Lane2:MCF-7 whole cell lysate(40ug)

Lane3:HCT116 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151