

CstF-50 (R3) polyclonal antibody

Catalog: BS2036

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Polyadenylation of mRNA precursors is a two-step reaction that requires multiple protein factors. The first step, endonucleolytic cleavage of polyadenylation substrates, requires CstF (cleavage stimulation factor), a heterotrimer that is composed of three distinct subunits of 77, 64 and 50 kDa. Heterotrimeric CstF recognizes GU and U-rich sequences located downstream of the polyadenylation site on RNA. The 50 kDa CstF subunit shares extensive homology with mammalian G protein beta-subunits and has a transducin repeat domain, which is a 44 amino acid-long sequence that is repeated seven times. CstF-50 interacts with the nuclear protein BARD1 (BRCA1-associated RING domain protein) and inhibits polyadenylation in vitro. CstF-50 may also be responsible for the interaction of the heterotrimeric CstF complex with other polyadenylation and 3'-end cleavage factors to form a stable complex on the pre-mRNA.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 50 kDa

Swiss-Prot:

Q05048

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

CstF-50 (R3) polyclonal antibody detects endogenous levels of CstF-50 protein

DATA:



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151