

CDK5R1 (Q31) polyclonal antibody

Catalog: BS2065

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Cyclin dependent kinase-5 (Cdk5), a key regulator of cell cycle progression, was originally isolated on the basis of its structural homology to Cdc2, a well characterized regulator of cell cycle progression. Although Cdk5 is expressed at the highest level in the brain of adult mice, intermediate levels in testis and low or undetectable levels in all other tissues, brain is the only tissue from which Cdk5 can be isolated as an active kinase. These findings may be explained by the cloning and characterization of a Cdk5 regulatory subunit, designated p35. p35 displays a neuronal cell-specific pattern of expression, physically associates with Cdk5 and activates Cdk5 enzymatic activity. p35 is also expressed in many tissues in a truncated form, designated p25.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 35 kDa

Swiss-Prot:

Q15078

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

CDK5R1 (Q31) polyclonal antibody detects endogenous levels of CDK5R1-p35 protein, and does not cross-react with CDK5R1-p25 protein.

DATA:



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151