

14-3-3 θ (Q67) polyclonal antibody

Catalog: BS2179

Host: Rabbit

Reactivity:

BackGround:

Protein interactions with 14-3-3 show distinct preferences for it's different isoforms and are regulated by phosphorylation of both 14-3-3 and the bound protein. 14-3-3 tau/theta is primarily expressed in T-cells although it occurs at lower levels in the brain, heart, pancreas placenta and kidneys. It plays a role in cell cycle progression via interaction with p27(Kip1). Along with COPI, the theta isoform binds the GB1 RSR sequence involved in protein trafficking. The isoform forms a complex with Hsp60 and cellular prion protein which may be involved in prion diseases. Elevated levels of 14-3-3 tau/theta are observed in amyotrophic lateral sclerosis. 14-3-3 tau/theta also BCR/Abl. bacterial effector binds protein Tir. yes-associated protein and the FSH receptor.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 28 kDa

Swiss-Prot:

P27348

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Specificity:

14-3-3 θ (Q67) polyclonal antibody detects endogenous levels of 14-3-3 protein theta/tau.

Bioworld Technology, Inc.


Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841

DATA:




Human

Western blot (WB) analysis of 14-3-3 θ (Q67) pAb at 1:500 dilution Lane1:K562 whole cell lysate(40ug) Lane2:HepG2 whole cell lysate(40ug) Lane3:A549 whole cell lysate(40ug) Lane4:HEK293T whole cell lysate(40ug) Lane5:U-87MG whole cell lysate(40ug)





Immunohistochemistry (IHC) analyzes of 14-3-3 θ (Q67) pAb in paraffin-embedded human breast carcinoma tissue at 1:50,showing cytoplasmic and nuclear staining.Negative control (the right)Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



Lot CJ36131

Immunohistochemistry (IHC) analyzes of 14-3-3 θ (Q67) pAb in paraffin-embedded human colon carcinoma tissue at 1:50, showing cytoplasmic and nuclear staining. Negative control (the right)Using PBS instead

Bioworld technology, co. Ltd.Add:No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.Email:info@biogot.comTel:0086-025-68037686Fax:0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology,Inc.

of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin

followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.		Biow	Bioworld technology, co. Ltd.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,	Add:	No 9, weidi road Qixia District Nanjing, 210046,	
	MN 55416,USA.		P. R. China.	
Email:	info@bioworlde.com	Email:	info@biogot.com	
Tel:	6123263284	Tel:	0086-025-68037686	
Fax:	6122933841	Fax:	0086-025-68035151	