

Ataxin-2 (F774) polyclonal antibody

Catalog: BS3811

Host: Rabbit

Reactivity: Human

BackGround:

Ataxin-2, the gene product of the human spinocerebellar ataxia type 2 (SCA2/ATXN2) gene, is a basic protein with two domains (Sm1 and Sm2) implicated in RNA splicing and protein interaction. Ataxin-2 interacts with a putative RNA-binding protein ataxin-2-binding-protein 1 (A2BP1), which is expressed in muscle and brain. Ataxin-2 is ubiquitously expressed with highest levels in the cytoplasm of Purkinje cells. Both A2BP1 and ataxin-2 are localized to the trans-Golgi network. Mice expressing ataxin-2 with polyglutamine show progressive functional deficits accompanied by loss of Purkinje cell dendritic arbor and eventually loss of Purkinje cells. In conclusion, expansion of ataxin-2 results in spinocerebellar ataxia type 2, which affects the cerebellum and other areas of the central nervous system.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 140 kDa

Swiss-Prot:

Q99700

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Ataxin-2 (F774) polyclonal antibody detects endogenous levels of Ataxin-2 protein.

DATA:

Immunohistochemistry (IHC) analyzes of Ataxin-2 (F774) pAb in paraffin-embedded human breast cancer tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: <u>info@bioworlde.com</u> Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151