

CXADR polyclonal antibody

Catalog: BS5638

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The coxsackie and adenovirus receptor (CAR) mediates viral infection by the binding of various adenoviruses through specific protein interactions. There is a high affinity between the viral knob domain and the extracellular amino-terminal domain (designated D1) of CAR. The D1 domain alone is sufficient for knob binding in transfected cells. Determining the specific interactions between CAR and adenoviruses is imperative in order to further develop gene therapy using viral hosts. CAR is expressed in many human and murine cell types. However, cells that express CAR at low levels are not efficiently infected by adenoviruses. Possible methods of avoiding this problem in certain cell types are by either supplementing CAR or modifying the Ad knob to bind to other receptors.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 46 kDa

Swiss-Prot:

P78310

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

ICC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

CXADR polyclonal antibody detects endogenous levels of CXADR protein.

DATA:



Western blot (WB) analysis of CXADR polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:NIH-3T3 cell lysate

Lane3:H9C2 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151