

CX3CL1 polyclonal antibody

Catalog: BS	0	U	0	1
-------------	---	---	---	---

Host:

: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Chemokines are members of a superfamily of inducible, secreted, proinflammatory cytokines. Members of the chemokine family exhibit 20-50%

9

homology in their predicted amino acid sequences and are divided into four

subfamilies. In the subfamily designated C-C or β , the first two cysteines are

adjacent. In the C-X-C or a subfamily, the first two of four cysteine residues

are separated by a single amino acid. C subfamily members, also designated

 γ chemokines, lack the first and third cysteine residues of the conserved

motif. Chemokines in these three subfamilies are small, secreted proteins

with molecular weights ranging from 7-15 kDa. Fractalkine, also designated

neurotactin, is the first characterized member of a fourth chemokine subfamily.

Fractalkine contains a novel C-X3-C motif in which the first two cysteines

are separated by three amino acid residues. Fractalkine mRNA has been

detected in brain and heart and is upregulated in microglia and endothelial

cells by inflammatory signals. The protein can exist both as a membranebound

form and as a chemotactic soluble form.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 42 kDa

Swiss-Prot:

P78423

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at $4 \,^{\circ}$ short term. Aliquot and store at $-20 \,^{\circ}$ long term. Avoid freeze-thaw cycles.

Specificity:

CX3CL1 polyclonal antibody detects endogenous levels of CX3CL1 protein.

DATA:



Western blot (WB) analysis of CX3CL1 polyclonal antibody at 1:500 dilution

Lane1:MCF-7 whole cell lysate

Lane2:NIH-3T3 whole cell lysate

Lane3:PC12 whole cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151