

Csk monoclonal antibody

Catalog:	MB0131
----------	--------

Host:

Mouse

Reactivity: Human, Rat

BackGround:

C terminal Src kinase (CSK) is a non receptor protein tyrosine kinase. One of its major functions is to specifically phosphorylate a conserved C terminal tyrosine of Src family kinases thereby down regulating their kinase activity. CSK is expressed ubiquitously as a protein of approximately 50 KDa in cytosol, but a fraction distributes to lipid rafts due to interaction with the lipid raft associated protein PAG (CSK binding protein).

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

Predicted band size:50KDa

Observed band size:50KDa

Swiss-Prot:

P41240

Purification&Purity:

The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000

ICC: 1:50

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at -20 $^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Specificity:

This antibody detects endogenous levels of Csk and does not cross-react with related proteins.

DATA:

_	JUNKO	්	
170KDa-			
130KDa-			
95KDa —			
72KDa -			
55KDa -			
43KDa _		-	Csk
34KDa -			

Western blot detection of CSK antibody in C6 and Jurkat cell lysates using CSK antibody (1:1000 diluted).



Immunocytochemistry staining of C6 cells fixed by anhydrous methanol for 2 h at -20° C and using anti-CSK antibody (dilution 1:50).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151