

Bioworld Technology, Inc.

Recombinant Human Tumor Necrosis Factor-alpha,His (rHuTNF-α-His)

Catalog Number: PR1126

Source: Escherichia coli.

Quantity:10µg/50µg/1.0mg

Description

Tumor necrosis factor alpha (TNF- α), also called cachectin, is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes endothelial cells, smooth muscle cells and some transformed cells. TNF- α occurs as a secreted, soluble form and as a membraneanchored form, both of which are biologically active. The naturally-occurring form of TNF- α is glycosylated, but nonglycosylated recombinant TNF- α has comparable biological activity. The biologically active native form of TNF- α is reportedly a trimer. Human and murine TNF-a show approximately 79% homology at the amino acid level and crossreactivity between the two species. Two types of receptors for TNF- α have been described and virtually all cell types studied show the presence of one or both of these receptor types.

Molecular Weight:

Approximately 17.5 kDa. a single, non-glycosylated, polypeptide chain containing 157 amino acids fragment (77-233) and having a molecular mass of 21.85 kDa with an amino-terminal hexahistidine tag.

Purity:

>95% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. The Specific Activity is $\geq 2.0 \times 107$ IU/mg as determined by the cytolysis of murine L929 cells in the presence of Actinomycin D

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.0.

AA Sequence:

M H H H H H H V R S S S R T P S D K P V A H V V A N P Q A E G Q L Q W L N R R A N A L L A N G V E L R D N Q L V V P S E G L Y L I Y S Q V L F K G Q G C P S T H V L L T H T I S R I A V S Y Q T K V N L L S A I K S P C Q RETPEGAEAKPWYEPIYLGGVFQLEKG DRLSAEINRPDYLDFAESGQVYFGIIAL

Endotoxin:

Less than 1EU/mg of rHuTNF- α -his as determined by LAL method.

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 10mM HAc to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

MADE IN CHINA Bioworld technology, co, Ltd. No 9, weidi road Qixia District Nanjing, 210046, P, R.China. Email: info@biogot.com Tel: 0086-025-86371664 Fax:0086-025-86213570

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841