

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Epigen, Human

Catalog Number: BK0031-50µg

Source: Escherichia coli.

Quantity: 50µg

Description:

Epigen is a cytokine belonging to the Epidermal Growth Factor (EGF) superfamily, which also includes Epiregulin, Amphiregulin, Neuregulin 2-β, and Transforming Growth Factor α. The precursor of Epigen produced in tissues has 154 amino acids, and shares the characteristics of other members of EGF superfamily, including 3 disulfide bonds formed by 6 cysteines. Epigen is present in testis, heart, and liver, and it binds to EGF receptors with a much lower binding affinity than EGF. However, Epigen is more mitogenic than EGF. Epigen achieves its strong mitogenic potency by suppressing ligand-induced receptor inactivation. Unlike EGF, Epigen can also bind to EGF receptors in low pH conditions, helping its recycling. Therefore Epigen has anomalous potency due to its prolonged presence. Recombinant human Epigen (rhEpigen) produced in E.coli is a single non-glycosylated polypeptide chain containing 73 amino acids. A fully biologically active molecule, rhEpigen has a molecular mass of 8.1 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Molecular Weight:

8.1 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% by SDS-PAGE analysis.

Biological Activity:

ED50 < 1 µg/mL, measured by a cell proliferation as-

say using 3T3 cells, corresponding to a specific activity of > 1 × 10³ units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

MAVTVTPPITAQQADNIEGPIALKF-
SHLCLEDHNSYCYINGACAFHHELEKAICRCFT-
GYTGERCEHLTLTSYA

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O at 100 µg/mL.

Storage:

Lyophilized recombinant human Epigen (rhEpigen) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rhEpigen remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.