PRODUCT DATA SHEET



Bioworld Technology,Inc.

Recombinant FGF-18, Mouse

Catalog Number: BK0038-10µg Source: Escherichia coli. Quantity: 10µg

Description:

Fibroblast Growth Factor 18 (FGF-18) is a pleiotropic cytokine belonging to the heparin-binding FGF family, which has 23 different members. Structurally, FGF-18 is closely related to FGF-8 and FGF-17. Like other FGFs, FGF-18 can bind to different FGF receptors in vivo. FGF-18 is expressed in various tissues and has multiple functions: during long bone growth, FGF-18 is expressed in perichondrium and developing joints, and regulates bone formation by inhibiting chondrocyte proliferation and differentiation: knock-out mice survive embryonic development, but exhibit skeletal abnormalities and die in the early neonatal period. FGF-18 also induces ectopic cartilage formation in the lung, and alters the morphology of the pulmonary mesenchyma. Recombinant mouse Fibroblast Growth Factor 18 (rmFGF-18) produced in E.coli is a single non-glycosylated polypeptide chain containing of 172 amino acids. A fully biologically active molecule, rmFGF-18 has a molecular mass of 20.1 kDa analyzed by non-reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Molecular Weight:

20.1 kDa, observed by non-reducing SDS-PAGE.

Purity

> 95% by SDS-PAGE and HPLC analyses.

Biological Activity:

ED50 < 0.5 μ g/mL, measured by a cell proliferation assay using 3T3 cells, corresponding to a specific activity of > 2× 10^3 units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

EENVDFRIHVE-

NQTRARDDVSRKQLRLYQLYSRTSGKHIQVLGR RISARGEDGDKYAQLL-

VETDTFGSQVRIKGKETEFYLCMNRKGKLVGKP DGTSKECVFIEKVLENNYTALMSAKYSGWY-VGFTKKGRPRKGPK-

TRENQQDVHFMKRYPKGQAELQKPFKYT-TVTKRSR

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH2O at 100 µg/mL.

Storage:

Lyophilized recombinant mouse Fibroblast Growth Factor 18 (rmFGF-18) remains stable up to 6 months at -80 $^{\circ}$ C from date of receipt. Upon reconstitution, rmFGF-18 remains stable up to 2 weeks at 4 $^{\circ}$ C or up to 3 months at -20 $^{\circ}$ C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.