

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant FGF-8a, Human

Catalog Number: BK0045-1mg

Source: Escherichia coli.

Quantity: 1mg

### Description:

Fibroblast Growth Factor 8a (FGF-8a) is a cytokine belonging to the heparin-binding FGF family, which has at least 23 members. FGF-8 has 8 different isoforms, named FGF-8a through FGF-8h. Different FGF-8 isoforms have different affinities to the receptors, and thus participate in different signaling cascade pathways. FGF-8 has very widespread expression during embryonic development, and is an organizer and inducer for gastrulation, somitogenesis, morphogenesis, and limb induction. However, FGF-8 is also a potential oncogene: in normal adult cells, FGF-8 has very low expression, but FGF-8 is highly expressed in cancer cells of breast, prostate, and ovarian tumors. FGF-8 promotes tumor angiogenesis by increasing neovascularization, and induces osteoblastic differentiation. Recombinant human Fibroblast Growth Factor 8a (rhFGF-8a) produced in E. coli is a single non-glycosylated polypeptide chain containing 183 amino acids. A fully biologically active molecule, rhFGF-8a has a molecular mass of 21.3 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

### Molecular Weight:

21.3 kDa, observed by reducing SDS-PAGE.

### Purity:

> 95% by SDS-PAGE analysis.

### Biological Activity:

ED50 < 500 ng/mL, measured by a cell proliferation assay using 3T3 cells in the presence of 1 µg/ml of heparin, corresponding to a specific activity of > 2 × 10<sup>3</sup> units/mg.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

MQHVVREQLSLVTDQLSRRLIR-  
TYQLYSRTSGKHFVQVLANK-  
RINAMAEDGDPFAKLIVETDTFGS-  
RVRVRGAETGLYICMNKKGKLIK-  
SNGKGDVCFTEIVLENNYALQNAKYEG-  
WYMAFTRKGRPRKGSKTRQHQREVHFMKRL-  
PRGHHTTEQSLRFEFLNY-  
PPFTRSLRGSQRTWAPEPR

### Endotoxin:

< 0.2 EU/µg, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O at 100 µg/mL.

### Storage:

Lyophilized recombinant human Fibroblast Growth Factor 8a (rhFGF-8a) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rhFGF-8a remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.