# PRODUCT DATA SHEET



Bioworld Technology,Inc.

# Recombinant FGF-9, Human

Catalog Number: BK0049-10µg Source: Escherichia coli. Quantity: 10µg

### **Description:**

Fibroblast Growth Factor-9 (FGF-9) is a heparin binding growth factor that belongs to the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-9 was isolated as a secreted factor that exhibits a growth-stimulating effect on cultured glial cells. In nervous system, this protein is produced mainly by neurons and may be important for glial cell development.Recombinant Human Fibroblast Growth Factor-9(rhFGF-9) produced in E.coli is a single non-glycosylated polypeptide chain containing 208 amino acids. A fully biologically active molecule, rhFGF-9 has a molecular mass of 23.4 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

### **Molecular Weight:**

23.4 kDa, observed by reducing SDS-PAGE.

#### **Purity:**

> 95% by SDS-PAGE and HPLC analyses.

### **Biological Activity:**

ED50 < 2.0 ng/ml , measured by a cell proliferation assay using 3T3 cells, corresponding to a specific activity of  $>5.0\times10^5$  units/mg.

#### **Physical Appearance:**

Sterile Filtered White lyophilized (freeze-dried) powder.

## **Formulation:**

Lyophilized after extensive dialysis against PBS.

### **AA Sequence:**

MAPLGEVGNYFGVQDAVPFGNVPVLPVD-SPVLLSDHLGQSEAGGLPRGPAVTDLDHLKGIL-RRRQLYCRTGFHLEIFPNGTIQGTRK-DHSRFGILEFISIAVGLVSIRGVDSGLYL-GMNEKGELYGSEKLTQECVFREQFEEN-WYNTYSSNLYKHVDTGRRYYVAL-NKDGTPREGTRTKRHQKFTH-FLPRPVDPDKVPELYKDILSQS

#### **Endotoxin:**

< 0.2 EU/µg, determined by LAL method.

#### **Reconstitution:**

Reconstituted in ddH2O at 100 µg/ml.

### **Storage:**

Lyophilized recombinant Human Fibroblast Growth Factor-9(rhFGF-9) remains stable up to 6 months at -80  $^{\circ}$ C from date of receipt. Upon reconstitution, rhFGF-9 should be stable up to 2 weeks at 4  $^{\circ}$ C or up to 3 months at -20  $^{\circ}$ C.

### **Usage:**

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.

Email: <u>info@bioworlde.com</u> Tel: 6123263284 Fax: 6122933841 Tel: 0086-025-86371664 Fax:0086-025-86213570