

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant GM-CSF, Human

Catalog Number: BK0066-10µg

Source: Escherichia coli.

Quantity: 10µg

Description:

Human Granulocyte Macrophage Colony Stimulating Factor (hGM-CSF), a hematopoietic growth factor, is mainly involved in granulopoiesis and monocytopenia. It is produced by T-cells and macrophages in response to antigens, and by endothelial cells and fibroblasts following induction of various cytokines [1]. A monomeric protein of 127 amino acids with six glycosylation sites and two intra disulfide bonds [2,3], glycosylated and non-glycosylated hGM-CSFs show similar biological activities [4]. Other than its connection to the growth and development of granulocytes and macrophages, it is also indispensable for the proliferation of erythroid and megakaryocytic cells [5]. Recombinant human Granulocyte Macrophage Colony Stimulating Factor (rhGM-CSF) produced in E.coli is a non-glycosylated polypeptide. A fully biologically active molecule, rhGM-CSF has a molecular mass of 14 kDa analyzed by SDS-PAGE and is obtained by proprietary refolding and chromatographic techniques at GenScript.

Molecular Weight:

14 kDa, observed by SDS-PAGE.

Purity:

>95 % as analyzed by SDS-PAGE.

Biological Activity:

ED50 < 0.5 ng/ml, measured by proliferation assay of TF-1 cells, corresponding to a specific activity of >2.0 x 10⁶ units/mg

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against 10 mM PB, pH7.0.

AA Sequence:

MAPARSPSPSTQPWEHVNAIQEARRLLNLSRD-
TAAEMNETVEVISEMFDLQEPTCLQTRLE-
LYKQGLRGSCLKLKGPLTMMASHYK-
QHCPTPETSCATQIIT-
FESFKENLKDPLLVIPFDCWEPVQE

Endotoxin:

<1 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O at 100 µg/ml.

Storage:

Lyophilized recombinant human Granulocyte Macrophage Colony Stimulating Factor (rhGM-CSF) remains stable up to 12 months at -80 °C from date of receipt. Upon reconstitution, rhGM-CSF should be stable up to 4 weeks at 4 °C or up to 6 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.