# PRODUCT DATA SHEET



Bioworld Technology,Inc.

# Recombinant HB-EGF, Human

Catalog Number: BK0073-10µg Source: Escherichia coli. Quantity: 10µg

# **Description:**

Proheparin-binding EGF-like growth factor (HB-EGF), also known as DTR, DTS and HEGFL, is a member of the EGF family of mitogens. It is expressed in macrophages, monocytes, endothelial cells and muscle cells. HB-EGF signals through the EGF receptor to stimulate the proliferation of smooth muscle cells, epithelial cells and keratinocytes. Compared to EGF, HB-EGF binds to the EGF receptor with a higher affinity and has been shown to bemore mitogenic, likely due to its ability to bind to heparin and heparin sulfate proteoglycans. HB-EGF has also been reported to act as a diphtheria toxin receptor, mediating endocytosis of the bound toxin. Heparin-binding EGF-like growth factor has been shown to interact with NRD1, Zinc finger and domain-containing protein BTB BAG1.Recombinant Human HB-EGF produced in E.coli cells is a polypeptide chain containing 86 amino acids. A fully biologically active molecule, rhHB-EGF has a molecular mass of 14 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

## **Molecular Weight:**

14 kDa, observed by reducing SDS-PAGE.

#### **Purity:**

> 95% as analyzed by SDS-PAGE.

## **Biological Activity:**

 $\mbox{ED50} < 0.75\mbox{ng/ml},$  measured in a cell proliferation assay using 3T3 cells.

# **Physical Appearance:**

Sterile Filtered White lyophilized (freeze-dried) powder.

## **Formulation:**

Lyophilized after extensive dialysis against PBS.

## **AA Sequence:**

GDLQEADLDLLRVTLSSKPQALATPNKEE-HGKRKKKGKGLGKKRDPCLRKYKDFCIH-GECKYVKELRAPSCICHPGYHGERCHGLSL

#### **Endotoxin:**

< 0.2 EU/µg, determined by LAL method.

## **Reconstitution:**

Reconstituted in ddH2O or PBS at 100 µg/ml.

#### **Storage:**

Lyophilized recombinant Human Proheparin-binding EGF-like Growth Factor (HB-EGF), remains stable up to 6 months at -80  $^{\circ}$ C from date of receipt. Upon reconstitution, human HB-EGF should be stable up to 1 week at 4  $^{\circ}$ C or up to 3 months at -20  $^{\circ}$ C.

#### Usage

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.