

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant IFN- $\gamma$ , Mouse

Catalog Number: BK0081-100 $\mu$ g

Source: Escherichia coli.

Quantity: 100 $\mu$ g

### Description:

Sharing 41% sequence identity with human Interferon gamma (hIFN- $\gamma$ ), mouse IFN gamma (mIFN- $\gamma$ ) is a macrophage-activating factor. The active form of IFN- $\gamma$  is an antiparallel dimer that sets off IFN- $\gamma$ /JAK/STAT pathway. IFN- $\gamma$  signaling does diverse biological functions primarily related to host defense and immune regulation, including antiviral and antibacterial defense, apoptosis, inflammation, and innate and acquired immunity. While IFN- $\gamma$ -induced inflammatory cascade summons a variety of immune-related cell types, such as macrophages, natural killer (NK) cells and cytotoxic T lymphocytes (CTLs), IFN- $\gamma$  is also implicated in resistance to NK cell and CTL responses and in immune escape in a variety of cancers. Recombinant mouse IFN gamma (rmIFN- $\gamma$ ) produced in E.coli is a non-glycosylated polypeptide chain of 134 amino acids. A fully biologically active molecule, rmIFN- $\gamma$  has a molecular mass of 15 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary refolding and chromatographic techniques at GenScript.

### Molecular Weight:

15 kDa, observed by reducing SDS-PAGE.

### Purity:

> 95% as analyzed by reducing SDS-PAGE.

### Biological Activity:

ED<sub>50</sub> < 0.15 ng/ml, measured by cytotoxicity assay using WEHI-279 cells.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

MHGTVIESLES LN NYFNSSGIDV E EKSLFLDI-  
WRNWQKDGDMKILQSQIISFYLR LFE-  
VLKDNQAISNNISVIESHLITTFNSN-  
SKAKKDAFMSIAKFEVNNPQVQRQAFNELIR-  
VVHQLPESSLRKRKR SRC

### Endotoxin:

< 1 EU/ $\mu$ g, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O or PBS at 100  $\mu$ g/ml.

### Storage:

Lyophilized recombinant mouse IFN gamma (rmIFN- $\gamma$ ) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rmIFN- $\gamma$  should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.