

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant IL-22, Mouse

Catalog Number: BK0101-50µg

Source: Escherichia coli.

Quantity: 50µg

### Description:

Interleukin-22 (IL-22) is a member of a group of cytokines called the IL-10 family which include IL-10, IL-19, IL-20, IL-24, and IL-26. IL-22 shares use of the IL-10R2 in cell signaling with other members of this family. IL-22 signals through a receptor system consisting of IL-10R-β/CRF2-4 and IL-22R, both of which are members of the class II cytokine-receptor family. IL-22 is produced by activated DCs and T cells and initiates an innate immune response against bacterial pathogens especially in epithelial cells such as those in the respiratory tract and gut. IL-22 along with IL-17 is rapidly produced by splenic LTi-like cells and can also be produced by Th17 cells, which plays a likely role in the coordinated response of both adaptive and innate immune systems. Recombinant Mouse IL-22 produced in E.coli is a single non-glycosylated polypeptide chain containing 147 amino acids. A fully biologically active molecule, rmIL-22 has a molecular mass of 16.8 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

### Molecular Weight:

16.8 kDa, observed by reducing SDS-PAGE

### Purity:

> 98% as analyzed by SDS-PAGE & HPLC.

### Biological Activity:

ED50 < 0.5 ng/ml. Measured by its ability to induce IL-10 secretion in COLO 205 (human colon carcinoma cells).

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

MLPVNTRCKLEVSNFQQPYYIVNRT-  
FMLAKEASLADNNTDVRLLIGEKLFGRG  
VSAKDQCYLMKQVLNFTLEDVLLPQS-  
DRFQPYMQEVVPFLTKLSNQLSSC  
HISGDDQNIQKNVRRLLKETVKKLGESEIKAI-  
GELDLLFMSLRNACV

### Endotoxin:

< 0.2 EU/µg, determined by LAL method.

### Reconstitution:

Reconstituted in ddH2O or PBS at 100 µg/ml.

### Storage:

Lyophilized recombinant Mouse IL-22 remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Mouse IL-22 should be stable up to 1 week at 4 °C or up to 3 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.