

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant IL-33, Human

Catalog Number: BK0104-10µg

Source: Escherichia coli.

Quantity: 10µg

Description:

Interleukin-33 (IL-33) is a proinflammatory cytokine that belongs to the IL-1 family. IL-33 is expressed in a variety of cells, including epithelial and endothelial cells, smooth muscle cells, macrophages and fibroblasts. The primary receptors for IL-33 are ST2 and IL-1 receptor accessory protein (IL-1RAcP), both of which belong to the IL-1 receptor family. IL-33 is localized to the nucleus of resting cells where it binds to chromatin in the H2A-H2B histone complex as a transcriptional suppressor. IL-33 is secreted by cells during injury which induces a T-helper 2 type inflammatory response. Evidence suggests IL-33 plays a role in autoimmune disease. IL-33's interaction with ST2 can drive allergic pathology and IL-33 has been reported to play a role in the development of rheumatoid arthritis and systemic lupus erythematosus. Recombinant human Interleukin-33 (rhIL-33) produced in E.coli is a single non-glycosylated polypeptide chain containing 159 amino acids. A fully biologically active molecule, rhIL-33 has a molecular mass of 18.0 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Molecular Weight:

18.0 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% by SDS-PAGE analysis.

Biological Activity:

ED50 <0.5ng/mL, measured by a cell proliferation assay using D10S cells, corresponding to a specific activity of $> 2 \times 10^6$ units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

SITGISPITEYLA SLSTYNDQSITFALEDESYE-
IYVEDLKKDEKKDKVLLSYYESQHP-
SNESGDGV D GKMLMVTL-
SPTKDFWLHANNKEHSVELHKCEKPLPDQAFFV
LHNMHNSNCVSFECKTDPGVFIGVKDNHLALIK-
VDSSENLCTENILFKLSET

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O at 100 µg/mL.

Storage:

Lyophilized recombinant human Interleukin-33 (rhIL-33) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rhIL-33 remains stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.