

## **Bioworld Technology,Inc.**

# **Recombinant MCP-1/CCL2, Human**

Catalog Number: BK0125-25µg

Source: Escherichia coli.

Quantity: 25µg

## **Description:**

Chemokine (C-C motif) ligand 2 (CCL2) is also referred to as monocyte chemotactic protein 1 (MCP1) and small inducible cytokine A2. CCL2 is a small cytokine that belongs to the CC chemokine family. CCL2 recruits monocytes, memory T cells, and dendritic cells to the sites of inflammation produced by either tissue injury or infection. Human CCL2 cDNA encodes a 99 amino acid (aa) precursor protein with a 23 aa signal peptide and a 76 aa mature protein. Human CCL2 shares 78-79% aa identity with canine, porcine and equine CCL2, while mouse and rat express a form of CCL2 that is extended by 49 aa and shares only ~56% aa identity within the common region.Recombinant Human MCP-1/CCL2 produced in E.coli is a single non-glycosylated polypeptide chain containing 76 amino acids. A fully biologically active molecule, rhMCP-1/CCL2 has a molecular mass of 8.7 kD analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

Molecular Weight:

8.7 kDa, observed by reducing SDS-PAGE.

#### **Purity:**

> 95% as analyzed by SDS-PAGE.

**Biological Activity:** 

The EC50 value of human MCP-1/CCL2on Ca<sup>2</sup>+ mobilization assay in CHO-K1/G $\alpha$ 15/hCCR2 cells (human G $\alpha$ 15 and human CCR2 stably expressed in CHO-K1 cells) is less than 1 $\mu$ g /ml.

## Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### **Formulation:**

Lyophilized after extensive dialysis against PBS.

#### **AA Sequence:**

QPDAINAPVTCCYNFTNRKISVQRLASYR-RITSSKCPKEAVIFK-TIVAKEICADPKQKWVQDSMDHLDKQTQTPKT

## **Endotoxin:**

 $< 0.2 \text{ EU/}\mu g$ , determined by LAL method.

#### **Reconstitution:**

Reconstituted in ddH2O or PBS at 100 µg/ml. **Storage:** 

Lyophilized recombinant Human MCP-1/CCL2 remains stable up to 6 months at -80  $^{\circ}$ C from date of receipt. Upon reconstitution, human MCP-1/CCL2 should be stable up to 1 week at 4  $^{\circ}$ C or up to 3 months at -20  $^{\circ}$ C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.