

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant OSM (227aa), Human

Catalog Number: BK0147-10µg

Source: Escherichia coli.

Quantity: 10µg

Description:

Oncostatin M (OSM) is a multifunctional cytokine, and belongs to Interleukin-6 (IL-6) subfamily, including IL-11, leukemia inhibitory factor (LIF), ciliary neurotropic factor, cardiotrophin-1, and novel neurotrophin-1. In vivo, OSM is secreted from activated T cells, monocytes, neutrophils, and endothelial cells. OSM is related to LIF, and share a receptor with LIF in human. Human OSM can bind to gp130 and recruit OSM Receptor β or LIF Receptor β to form a ternary complex. OSM stimulates the growth of different types of cells, including megakaryocytes, fibroblasts, vascular endothelial cells, and T cells. On the other hand, OSM inhibits the proliferation of several cancer cell lines, such as solid tissue tumor cells, lung cancer cells, melanoma cells, and breast cancer cells. Recombinant human Oncostatin M (rhOSM) produced in E.coli is a single non-glycosylated polypeptide chain containing 228 amino acids. A fully biologically active molecule, rhOSM has a molecular mass of 25.9 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Molecular Weight:

25.9 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% by SDS-PAGE analysis.

Biological Activity:

ED50 < 10 ng/mL, measured by a cell proliferation assay using TF-1 cells, corresponding to a specific activity of > 1×10^5 units/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

MAAIG-
SCSKEYRVLLGQLQKQTDLMQDTSRLLDPIYIRIQ
GLDVPKLRHCRERPGAFPSEETLR-
GLGRRGFLQTLNATLGCVLHRLADLEQRL-
PKAQDLERSGLNIEDLEKLMARPNILGLRN-
NIYCMAQLLDNSDTAEPT-
KAGRGASQPPTPTPASDAFQRKLEGCRFLHGY-
HRFMHSVGRVFSKWGESPNRSRRHSPHQAL-
RKGVRTRPSRKGKRLMTRGQLPR

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O at 100 µg/mL.

Storage:

Lyophilized recombinant human Oncostatin M (rhOSM) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rhOSM should be stable up to 2 weeks at 4 °C or up to 3 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.