

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant HCC-4/CCL16, Human(CHO-expressed)

Catalog Number: BK0219-10µg

Source: CHO

Quantity: 10µg

### Description:

Human HCC 4, also named NCC 4 and Chemokine (C-C motif) ligand 16 (CCL16) is a small cytokine belonging to the CC chemokine family that is known under several pseudonyms, including Liver-expressed chemokine (LEC) and Monotactin-1 (MTN-1). It can signal through the CCR8 and CCR1 receptors, and it is chemotactic towards monocytes and lymphocytes but not neutrophils. HCC-4 is expressed weakly by some lymphocytes, including NK cells, T cells, and some T cell clones. The expression of HCC-4 in monocytes is greatly up-regulated in the presence of IL-10. HCC-4 induces a calcium flux in thp-1 cells that are desensitized prior to the expression of RANTES. Recombinant human HCC-4/CCL16 produced in CHO cells is a single non-glycosylated polypeptide chain containing 97 amino acids. A fully biologically active molecule, rhHCC-4/CCL16 has a molecular mass of 12 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

### Molecular Weight:

12 kDa, observed by reducing SDS-PAGE.

### Purity:

> 98% as analyzed by SDS-PAGE.

### Biological Activity:

The EC<sub>50</sub> value of human HCC-4/CCL16 on Ca<sup>2+</sup> mobilization assay in CHO-K1/Ga15/hCCR1 cells (human Ga15 and human CCR1 stably expressed in CHO-K1 cells) is less than 1.5 µg/ml.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

QPKVPEWVNTTPSTCCLKYYEKVLPRLVVGYRK  
ALNCHLPAIIFVTKRNREVCTNPNDWVQEY-  
IKDPNLPLLPTRNLSTVKIITAKNGQPQLLSQ

### Endotoxin:

< 0.2 EU/µg, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O or PBS at 100 µg/ml.

### Storage:

Lyophilized recombinant human HCC-4/CCL16 remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, human HCC-4/CCL16 should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.