

# PRODUCT DATA SHEET



Bioworld Technology, Inc.

## Recombinant IL-10, Mouse(CHO-expressed)

Catalog Number: BK0226-50µg

Source: CHO

Quantity: 50µg

### Description:

Interleukin-10 (IL-10), initially known as Cytokine Synthesis Inhibitory Factor (CSIF), belongs to the IL-10 family and shares more than 80% sequence homology with the Epstein-Barr Virus protein BCRFI. It is produced by many immune cells, such as T-cells, macrophages, mast cells and dendritic cells. It is usually secreted as a homodimer and, upon binding to its receptor, inhibits the synthesis of a number of cytokines, including IFN-gamma, IL-2, IL-3, TNF and GM-CSF, by activated macrophages and Th2 cells. It also displays the ability to suppress Antigen-Presenting Cell (APC) function. The net effect of Interleukin-10 appears to be inhibitory; however, stimulatory effects, such as stimulation of B cell maturation and antibody production, are also reported.

### Molecular Weight:

21-23 kDa, observed by reducing SDS-PAGE.

### Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

### Biological Activity:

ED50 <0.2 ng/ml, measured in a cell proliferation assay using MC/9 cells.

### Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

### Formulation:

Lyophilized after extensive dialysis against PBS.

### AA Sequence:

SRGQYSREDNNC-  
THFPVQGSHMLLELRTAFSQQVKTFFQTKDQLD-  
NILLTDSLMLQDFKGYLGCQALSEMIQFYL-  
VEVMPQAEKHGPEIKEHLNSLGEKLTLM-  
RLRRCHRFLPCENKSKAVEQVKS-  
DFNKLQDQGVYKAMNEFD-  
IFINCIEAYMMIKMKS

### Endotoxin:

< 0.2 EU/µg, determined by LAL method.

### Reconstitution:

Reconstituted in ddH<sub>2</sub>O or PBS at 100 µg/ml.

### Storage:

Lyophilized recombinant murine Interleukin-10 remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, murine Interleukin-10 should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.