

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant IL-3, His, Rat

Catalog Number: BK0242-1mg

Source: CHO

Quantity: 1mg

Description:

Interleukin-3 (IL-3), also known as MCGF, Multi-CSF, HCGF and P-cell stimulation factor, belongs to the α -helix family of hematopoietic cytokines. It is produced by activated T-cells, mast cells and natural killer cells. IL-3 binds to the IL-3 receptor alpha subunit and recruits the signal-transducing common beta chain. IL-3 induced signal transduction results in the survival, differentiation and proliferation of a variety of immune cells, such as macrophages, neutrophils, megakaryocytes, mast cells and hematopoietic stem cells. IL-3 often acts synergistically with other cytokines, such as IL-7, EPO, GM-CSF and IL-6, to exert its stimulative function.

Molecular Weight:

22-34 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE.

Biological Activity:

ED50 < 8 ng/ml, measured in a cell proliferation assay using NF S-60 cells.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

ISDRGSDAHLLRLTD-
CRTIALEILVKLPYPQVSGLNNSDDKANLRNSTL
RRVNLDEFLKSQEEFDSQDITDIK-
SKLQKLKCCIPAAASDSVLPGVYN-
KDLDDFKKKLRFYVIHLKDLQPVSVSRPPQPTSS
SDNFRPMTVECHHHHHH

Endotoxin:

< 0.2 EU/ μ g, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 μ g/ml.

Storage:

Lyophilized recombinant rat Interleukin-3 (IL-3), His remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, rat Interleukin-3 (IL-3), His should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.