

Bioworld Technology,Inc.

Recombinant IL-6, Human(CHO-expressed)

Catalog Number: BK0253-50µg

Source: CHO

Quantity: 50µg

Description:

Interleukin-6 (IL-6), also known as BSF-2, CDF and IFNB2, is a pleiotropic cytokine that acts in both pro-inflammatory and anti-inflammatory responses. It is produced mainly by T cells, macrophages, monocytes, endothelial cells and muscle cells. IL-6 binds to IL-6 receptor (IL-6R) to trigger the association of IL-6R with gp130, inducing signal transduction through JAKs and STATs. The biological functions of IL-6 are diverse. It stimulates B cell differentiation and antibody production, myeloma and plasmacytoma growth, as well as nerve cell differentiation. It also acts as a myokine, produced by muscle cells in response to muscle contraction, to be released to the blood stream to help break down fats and improve insulin resistance.

Molecular Weight:

21-23 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE.

Biological Activity:

ED50 < 0.06 ng/ml, measured in a cell proliferation assay using 7TD1 cells.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

PVPPGEDSKDVAAPHRQPLTSSERIDKQIRYILD-GISALRKETCNKSNMCESS-KEALAENNLNLPKMAEKDGCFQSGFNEETCLVK IITGLLEFEVYLEYLQNRFESSEEQARAVQM-STKVLIQFLQKKAKNLDAITTPDPTT-NASLLTKLQAQNQWLQDMTTHLILRS-FKEFLQSSLRALRQM

Endotoxin:

 $< 0.2 \text{ EU/}\mu g$, determined by LAL method.

Reconstitution:

Reconstituted in ddH2O or PBS at 100 µg/ml. **Storage:**

Lyophilized recombinant Human Interleukin-6 remains stable up to 6 months at -80 $^{\circ}$ C from date of receipt. Upon reconstitution, Human Interleukin-6 should be stable up to 1 week at 4 $^{\circ}$ C or up to 2 months at -20 $^{\circ}$ C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.