

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant MIP-3 α /CCL20, Human(CHO-expressed)

Catalog Number: BK0272-5 μ g

Source: CHO

Quantity: 5 μ g

Description:

Chemokine (C-C motif) ligand 20 (CCL20) also known as liver activation regulated chemokine (LARC) or Macrophage Inflammatory Protein-3 (MIP3 α) is a small cytokine belonging to the CC chemokine family. It is strongly chemotactic for lymphocytes and weakly attracts neutrophils. Additionally, it promotes the adhesion of memory CD4⁺ T cells and inhibits colony formation of bone marrow myeloid immature progenitors. Recombinant human MIP-3 α /CCL20 produced in CHO cells is a polypeptide chain containing 70 amino acids. A fully biologically active molecule, rhMIP-3 α /CCL20 has a molecular mass of 8kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

Molecular Weight:

8 kDa, observed by reducing SDS-PAGE.

Purity:

> 98% as analyzed by SDS-PAGE.

Biological Activity:

The EC₅₀ value of human MIP-3 α /CCL20 on Ca²⁺ mobilization assay in CHO-K1/Ga15/hCCR6 cells (human Ga15 and human CCR6 stably expressed in CHO-K1 cells) is less than 200 ng/ml.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

ASNFDCCCLGYT-
DRILHPKFIVGFTRQLANEGCDINAIIFHT-
KKKLSVCANPKQTWVKYIVRLLSKKVKNM

Endotoxin:

< 0.2 EU/ μ g, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 μ g/ml.

Storage:

Lyophilized recombinant Human MIP-3 α /CCL20 remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Human MIP-3 α /CCL20 should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.