

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant TGF- α , Human

Catalog Number: BK0285-25 μ g

Source: CHO

Quantity: 25 μ g

Description:

Protransforming Growth Factor-alpha (TGF-alpha), also known as sarcoma growth factor, TGF-type I and ETGF, is a member of the EGF family of cytokines. It is expressed in monocytes, brain cells, keratinocytes and various tumor cells. ProTGF-alpha signals through EGFR and acts synergistically with TGF-beta to promote the proliferation of a wide range of epidermal and epithelial cells. It may function as either a membrane-bound ligand or a soluble ligand. Membrane-bound proTGF-alpha plays a role in cell-cell adhesion and juxtacrine stimulation of adjacent cells. The soluble form of the cytokine is released from the membrane-bound form by proteolytic cleavage and acts as a mitogen for cell proliferation.

Molecular Weight:

8-10 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

Biological Activity:

ED50 < 0.4 ng/ml, measured in a cell proliferation assay using 3T3 cells.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) pow-

der.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

VVSHFNDCPDSHTQFCFHGTCRFLVQEDK-
PACVCHSGYVGARCEHADLLA

Endotoxin:

< 0.2 EU/ μ g, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 μ g/ml.

Storage:

Lyophilized recombinant Human TGF-alpha remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Human TGF-alpha Receptor should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.