

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant TNF R I, Human

Catalog Number: BK0287-10 μ g

Source: CHO

Quantity: 10 μ g

Description:

TNF Receptor Type I, is also known as TNF R-p55/p60 and TNFRSF1A. It is a type I transmembrane protein member of the TNF receptor superfamily. It is expressed in most cell types. Binding of either TNF- α or TNF- β to TNF-R1 initiates a signal transduction pathway that results in the activation of the transcription factor NF- κ B, whose target genes are involved in the regulation of inflammatory responses, and, in certain cells, induce apoptosis. TNF-R1 is essential for proper development of lymph node germinal centers and Peyer's patches and for combating intracellular pathogens such as Listeria. It is stored in the Golgi and translocates to the cell surface following proinflammatory stimuli.

Molecular Weight:

28~35 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE.

Biological Activity:

ED50 < 50 ng/ml, measured in a cell proliferation assay using 929 cells in the presence of 1 ng/ml human TNF- α .

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

DSVCPQGKYIHPQNNICCTK-
CHKGTYLYNDCPGPGQDTDCRECEGS-
FTASENHLRHCLSCSKCRKEMGQVEISS-
CTVDRDTCVCGCRKNQYRHYWSEN-
LFQCFNCSLCLNGTVHLSCQEKQNTVCTCHAG-
FFLRENECVSCSNCKKSLECTKLCLPQIEN

Endotoxin:

< 0.2 EU/ μ g, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 μ g/ml.

Storage:

Lyophilized recombinant Human TNF Receptor Type I remains stable up to 6 months at -80 $^{\circ}$ C from date of receipt. Upon reconstitution, Human TNF Receptor Type I should be stable up to 1 week at 4 $^{\circ}$ C or up to 2 months at -20 $^{\circ}$ C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.