

## **Bioworld Technology,Inc.**

# **Recombinant Amphiregulin, Human**

Catalog Number: BK0297-10µg

Source: HEK 293

Quantity: 10µg

### **Description:**

Amphiregulin is a member of the EGF family of cytokines, which comprises at least ten proteins including EGF, TGF- $\alpha$ , HB-EGF, Epiregulin, Tomoregulin, Neuregulins and the various heregulins. Through the EGF/TGF- $\alpha$  receptor, it stimulates growth of keratinocytes, epithelial cells and some fibroblasts. Amphiregulin also inhibits the growth of certain carcinoma cell lines. Synthesized as a transmembrane protein, Amphiregulin's extracellular domain is proteolytically processed to release the mature protein.

### Molecular Weight:

15~20 kDa, observed by reducing SDS-PAGE.

### **Purity:**

> 95% as analyzed by SDS-PAGE.

**Biological Activity:** 

ED50 < 0.2 ng/ml, measured in a cell proliferation assay using 3T3 cells.

**Physical Appearance:** 

Sterile Filtered White lyophilized (freeze-dried) powder.

**Formulation:** 

Lyophilized after extensive dialysis against PBS.

#### **AA Sequence:**

SVRVEQVVKPPQNKTESENTSDKPKRK-KKGGKNGKNRRNRKKKNPCNAEFQNFCIH-GECKYIEH-LEAVTCKCQQEYFGERCGEKSMKTHSMIDSSLS K

# **Endotoxin:**

 $< 0.2 \text{ EU/}\mu g$ , determined by LAL method.

**Reconstitution:** 

Reconstituted in ddH2O or PBS at 100 µg/ml.

### **Storage:**

Lyophilized recombinant Human Amphiregulin remains stable up to 6 months at -80  $^{\circ}$ C from date of receipt. Upon reconstitution, Human Amphiregulin should be stable up to 1 week at 4  $^{\circ}$ C or up to 2 months at -20  $^{\circ}$ C.

#### Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.