

Bioworld Technology,Inc.

Recombinant IL-1 RA, Human(HEK 293-expressed)

Catalog Number: BK0310-50µg

Source: HEK 293

Quantity: 50µg

Description:

IL-1 Receptor Antagonist, also known as IL-1RA, ICIL-1RA, IRAP and IL-1RN, is a member of the interleukin 1 cytokine family. It is expressed by monocytes, neutrophils, macrophages, epithelial cells and fibroblasts. IL-1RA inhibits the activity of both IL-1alpha and IL-1beta, and modulates a variety of IL-1 related immune and inflammatory responses. It inhibits the activity of IL-1 by binding to the receptor IL-1R1 and preventing its association with the coreceptor IL-1RAP for signaling. Clinical studies are being conducted to investigate the use of IL-1RA in the treatment of sepsis, rheumatoid arthritis and chronic myelogenous leukemia.

Molecular Weight:

18-23 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

Biological Activity:

 $ED50 < 0.1~\mu g$ /ml, measured in a neutralization assay using D10S cells in the presence of 50pg/ml Human IL-1a.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

RPSGRKSSKMQAFRIWDVNQKTFYLRNNQL-VAGYLQGPNVNLEEK-IDVVPIEPHALFLGIHGGKMCLSCVKSGDETRLQ LEAVNITDLSENRKQDKRFAF-IRSDSGPTTSFESAACPGW-FLCTAMEADQPVSLTNMPDEGVMVTKFYFQEDE **Endotoxin:** < 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH2O or PBS at 100 μ g/ml.

Storage:

Lyophilized recombinant human IL-1 Receptor Antagonist remains stable up to 6 months at -80 $^{\circ}$ C from date of receipt. Upon reconstitution, Human IL-1 Receptor Antagonist should be stable up to 1 week at 4 $^{\circ}$ C or up to 2 months at -20 $^{\circ}$ C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.