

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant IL-11, Mouse(HEK 293-expressed)

Catalog Number: BK0311-50µg

Source: HEK 293

Quantity: 50µg

Description:

Interleukin-11 (IL-11) is a pleiotropic cytokine that was originally detected in the conditioned medium of an IL-1 α -stimulated primate bone marrow stromal cell line (PU-34) as a mitogen for the IL-6-responsive mouse plasmacytoma cell line T11. IL-11 contains no cysteine residues or potential glycosylation sites. IL-11 has multiple effects on both hematopoietic and non-hematopoietic cells. Many of the biological effects described for IL-11 overlap those for IL-6. In vitro, IL-11 can synergize with IL-3, IL-4 and SCF to shorten the G0 period of early hematopoietic progenitors. IL-11 also enhances the IL-3-dependent megakaryocyte colony formation. IL-11 has been found to stimulate the T cell dependent development of specific immunoglobulin-secreting B cell.

Molecular Weight:

~21 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

Biological Activity:

ED50 < 3 ng/ml, measured in a cell proliferation assay using T11 cells.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

GPPAGSPRVSSDPRADLDSAVLLTRSLLD-
TRQLAAQMRDKFPADGDHSLDSLPTLAMSAG-
TLGSLQLPGVLTRLRVDLMSYLRHVQWL-
RRAGGPSLKTLEPEL-
GALQARLERLLRRLQLLMSRLALPQAAPDQPVIP
LGPPASAWGSIRAAHAILGGLHLTLDWAVR-
GLLLKTRL

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH2O or PBS at 100 µg/ml.

Storage:

Lyophilized recombinant Murine Interleukin-11 (IL-11) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Murine Interleukin-11 (IL-11) should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.