

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant IL-22, Human(HEK 293-expressed)

Catalog Number: BK0315-10µg

Source: HEK 293

Quantity: 10µg

Description:

Interleukin-22 (IL-22) is a member of the IL-10 family of regulatory cytokines which includes IL-10, IL-19, IL-20, IL-22, IL-24 and IL-26. Members of this family share partial homology in their amino acid sequences, but they are dissimilar in their biological functions. Produced by T lymphocytes, IL-22 inhibits IL-4 production by Th2 cells, and induces acute phase reactants in the liver and pancreas. IL-22 signals through a receptor system consisting of IL-10R-β/CRF2-4 and IL-22R, both of which are members of the class II cytokine-receptor family.

Molecular Weight:

16~28 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE.

Biological Activity:

Active, measured by its ability to activate STAT following receptor ligand interaction.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

APISSHCRLDKSNFQQPYITNRT-
FMLAKEASLADNNTDVRLLIGEKL-
HGVSMSERCYLMKQVLNFTLEEVLPQS-
DRFQPYMQEVVPLARLS-
NRLSTCHIEGDDLHIQRNVQKLKDTVKKLGESG
EIKAI GELDLLFMSLRNACI

Endotoxin:

< 0.2 EU/µg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 µg/ml.

Storage:

Lyophilized recombinant Human Interleukin-22 (IL-22) remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Human Interleukin-22 (IL-22) should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.