

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### HDAC1 (E468) Peptide

Cat No.: BS1160P

#### Background

Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, PCAF (for p300/CBP-associated factor), p300/CBP and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated mammalian RPD3) and HDAC3, all of which are related to the yeast transcriptional regulator Rpd3p, have been identified as histone deacetylases.

#### Swiss-Prot

Q13547

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS1160 HDAC1 (E468) pAb.

#### Purification & Purity

Synthetic peptide HDAC1 (E468). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.