

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



HDAC 9 (P1047) Peptide

Cat No.: BS1168P

Background

Acetylation of lysine residues in the amino terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Several mammalian proteins function as nuclear histone acetylases, including GCN5, PCAF (p300/CBP-associated factor), p300/CBP, HAT1 and the TFIID subunit TAF II p250. Conversely, the deacetylation of histones is associated with transcriptional silencing. The histone deacetylases (HDAC) include HDAC1-9. HDAC9 and HDAC9a are two alternatively spliced isoforms of HDAC9. HDAC9a is 132 amino acids shorter than HDAC9, but both isoforms contain the HDAC catalytic domain, remain capable of deacetylase activity and repress myocyte enhancer-binding factor 2-mediated transcription.

Swiss-Prot

Q9UKV0

Applications

Blocking

Specificity

This peptide can be used with studies using BS1168 HDAC 9 (P1047) pAb.

Purification & Purity

Synthetic peptide HDAC 9 (P1047). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.