

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



Integrin $\beta 3$ (T767) Peptide

Cat No.: BS1187P

Background

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16α and 8β subunits heterodimerize to produce more than 20 different receptors. Most Integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, Collagen and Vitronectin. Certain Integrins can also bind to soluble ligands such as Fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster Integrins by binding to adjacent Integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of Integrin-mediated responses.

Swiss-Prot

P05106

Applications

Blocking

Specificity

This peptide can be used with studies using BS1187 Integrin $\beta 3$ (T767) pAb.

Purification & Purity

Synthetic peptide Integrin $\beta 3$ (T767). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.