Bioworld Technology CO., Ltd.



p53 (P27) Peptide

Cat No.: BS1275P

Background

p53, a DNA-binding, oligomerization domain and transcription activation domain-containing tumor suppressor, upregulates growth arrest and apoptosisrelated genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus, yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 poly-ubiquitinates p53 for proteasome targeting. p53 fluctuates between latent and active (DNA-binding) conformations and is differentially activated through posttranslational modifications including phosphorylation and acetylation.

Swiss-Prot

P04637

Applications

Blocking

Specificity

This peptide can be used with studies using BS1275 p53 (P27) pAb.

Purification & Purity

Synthetic peptide p53 (P27). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.