Bioworld Technology CO., Ltd.



PR (V184) Peptide

Cat No.: BS1299P

Background

The effects of progesterone are mediated by two functionally different isoforms of the progesterone receptor, PR-A and PR-B, which are transcribed from distinct, estrogen inducible promoters within a single copy of the PR gene. The PR-A and PR-B proteins are 90 kDa and 118 kDa respectively; the first 164 amino acids of PR-B are absent in PR-A. Progesterone bound PR-A and PR-B have different transcription activation properties. Specifically, PR-B functions as a transcriptional activator in most cell and promoter contexts, while PR-A is transcriptionally inactive and functions as a strong ligand dependent transdominant repressor of steroid hormone receptor transcriptiona activity. An inhibitory domain (ID), which maps to the amino terminus of the receptor, exists within both PR isoforms.

Swiss-Prot

P06401

Applications

Blocking

Specificity

This peptide can be used with studies using BS1299 PR (V184) pAb.

Purification & Purity

Synthetic peptide PR (V184). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.