

## ACC $\alpha$ (H74) polyclonal antibody

Catalog: BS1378

Host: Rabbit

Reactivity: Human, Mouse, Rat

### Background:

Acetyl-CoA carboxylase (ACC) is a complex multifunctional enzyme system which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, the ratelimiting step in fatty acid synthesis. Exercise diminishes the activity of acetyl-CoA carboxylase in human muscle. ACC $\alpha$  (ACC1) is the rate-limiting enzyme in the biogenesis of long-chain fatty acids, and ACC $\beta$  (ACC2) may control mitochondrial fatty acid oxidation. These two isoforms of ACC control the amount of fatty acids in the cells. The catalytic function of ACC $\alpha$  is regulated by phosphorylation (inactive) and dephosphorylation (active) of targeted Serine residues and by allosteric transformation by citrate or palmitoyl-CoA, which serve as the short-term regulatory mechanism of the enzyme. The gene encoding ACC $\alpha$ , which maps to human chromosome 17, encodes the 265 kDa  $\alpha$  form of ACC, which is the major ACC in lipogenic tissues. The catalytic core of ACC $\beta$  is homologous to that of ACC $\alpha$  except for an additional peptide of about 150 amino acids at the N-terminus.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 265 kDa

### Swiss-Prot:

Q13085

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

### Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

### Specificity:

ACC $\alpha$  (H74) polyclonal antibody detects endogenous levels of ACC $\alpha$  protein.


### DATA:



Western blot (WB) analysis of ACC $\alpha$  (H74) polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:NIH-3T3 cell lysate



BS1378  
Lot CA36131

Immunohistochemistry (IHC) analyzes of ACC $\alpha$  (H74) pAb in paraffin-embedded human liver carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151



## PRODUCT DATA SHEET

Bioworld Technology, Inc.

---

---

### **Bioworld Technology, Inc.**

**Add:** 1660 South Highway 100, Suite 500 St. Louis Park,  
MN 55416, USA.

**Email:** [info@bioworld.com](mailto:info@bioworld.com)

**Tel:** 6123263284

**Fax:** 6122933841

### **Bioworld technology, co. Ltd.**

**Add:** No 9, weidi road Qixia District Nanjing, 210046,  
P. R. China.

**Email:** [info@biogot.com](mailto:info@biogot.com)

**Tel:** 0086-025-68037686

**Fax:** 0086-025-68035151