

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### PCAF (F820) Peptide

Cat No.: BS1417P

#### Background

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, PCAF (for p300/CBP-associated factor), p300/CBP and the TFIID subunit TAFII p250. Mammalian HDAC1 (also designated HD1) and HDAC2 (also designated mammalian RPD3), both of which are related to the yeast transcriptional regulator Rpd3p, have been identified as histone deacetylases.

#### Swiss-Prot

Q92831

#### Applications

#### Blocking

#### Specificity

This peptide can be used with studies using BS1417 PCAF (F820) pAb.

#### Purification & Purity

Synthetic peptide PCAF (F820). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.  
1660 South Highway 100, Suite 500 St. Louis Park, MN  
55416, USA. Email: [info@bioworld.com](mailto:info@bioworld.com)  
Tel: 6123263284 Fax: 6122933841

Bioworld technology, co, Ltd.  
No 9, weidi road Qixia District Nanjing, 210046,  
P, R.China. Email: [info@biogot.com](mailto:info@biogot.com)  
Tel: +86-025-68037686 Fax: +86-025-68035151