

## TNF-R1 (A405) polyclonal antibody

Catalog: BS1478

Host: Rabbit

Reactivity: Human, Mouse, Rat

### Background:

Tumor Necrosis Factor (TNF) is a cytokine whose function is mediated through two distinct cell surface receptors (TNF Receptor I and TNF Receptor II) that are included in the TNF Receptor superfamily along with FAS antigen and CD40. TNF Receptors I and II are 55 and 75 kDa members, respectively, of a family of cell surface molecules including nerve growth factor receptor, Fas/Apo1, CD30, OX40, and 41BB, which are characterized by cysteine rich motifs in the extracellular domain. While TNF Receptor I and TNF Receptor II share 28% sequence homology in the extracellular domains, their intracellular domains lack sequence homology, suggesting that they differ in their internal signal transduction pathways. TNF Receptor I contains an approximately 80 amino acid death domain near its carboxy terminus capable of transmitting an apoptotic signal through its interaction with TRADD (TNF Receptor I associated death domain protein), and subsequent interactions with FADD.

### Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

### Molecular Weight:

~ 50 kDa

### Swiss-Prot:

P19438

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

### Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

### Specificity:

TNF Receptor-1 (A405) polyclonal antibody detects endogenous levels of TNF Receptor-1 protein.

### DATA:




Western blot (WB) analysis of TNF-R1 (A405) polyclonal antibody at 1:500 dilution

Lane1:Hela whole cell lysate(40ug)

Lane2:Hela treated with H2O2 (100nmol/ml, 25min, phostop) whole cell lysate (40ug)

Lane3:Hela treated with H2O2 (100nmol/ml, 25min, λ-ppase) whole cell lysate (40ug)



Immunohistochemistry (IHC) analyzes of TNF-R1 (A405) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151