

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



TNF-R1 (A405) Peptide

Cat No.: BS1478P

Background

Tumor Necrosis Factor (TNF) is a cytokine whose function is mediated through two distinct cell surface receptors (TNF Receptor I and TNF Receptor II) that are included in the TNF Receptor superfamily along with FAS antigen and CD40. TNF Receptors I and II are 55 and 75 kDa members, respectively, of a family of cell surface molecules including nerve growth factor receptor, Fas/Apo1, CD30, OX40, and 41BB, which are characterized by cysteine rich motifs in the extracellular domain. While TNF Receptor I and TNF Receptor II share 28% sequence homology in the extracellular domains, their intracellular domains lack sequence homology, suggesting that they differ in their internal signal transduction pathways. TNF Receptor I contains an approximately 80 amino acid death domain near its carboxy terminus capable of transmitting an apoptotic signal through its interaction with TRADD (TNF Receptor I associated death domain protein), and subsequent interactions with FADD.

Swiss-Prot

P19438

Applications

Blocking

Specificity

This peptide can be used with studies using BS1478 TNF-R1 (A405) pAb.

Purification & Purity

Synthetic peptide TNF-R1 (A405). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.