

ETK (K35) polyclonal antibody

Catalog: BS1536

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

ETK, also known as BMX, is a member of the Bruton's tyrosine kinase (Btk) family. It is expressed in a variety of hematopoietic, epithelial and endothelial cells. Etk, like other Btk family members, contains a pleckstrin homology (PH) domain and Src homology SH3 and SH2 domains. It participates in multiple signal transduction pathways. Phosphorylation of Tyr566 by Src kinase is required for activation of Etk in vivo. In endothelial and epithelial cells, Etk is regulated by FAK through phosphorylation at Tyr40.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 78 kDa

Swiss-Prot:

P51813

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


ETK (K35) polyclonal antibody detects endogenous levels of ETK protein.

DATA:



BS1536
Lot CJ36131

Immunohistochemistry (IHC) analyzes of ETK (K35) pAb in paraffin-embedded human liver carcinoma tissue at 1:50, showing cytoplasmic and cell membrane staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS1536
Lot CJ36131

Immunohistochemistry (IHC) analyzes of ETK (K35) pAb in paraffin-embedded human rectum carcinoma tissue at 1:50, showing cytoplasmic and cell membrane staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151