

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### PEA-15 (D110) Peptide

Cat No.: BS1573P

#### Background

PEA-15 (phosphoprotein enriched in astrocytes) exists in an non-phosphorylated form (N), and two phosphorylated forms, Pa and Pb. PEA-15 is an endogenous substrate for PKC, which mediates the transition from Pa to Pb. The level of PEA-15 phosphorylation changes upon depolymerization or stabilization of tubulins, indicating that PEA-15 co-localizes with microtubules. The first 80 amino acids of PEA-15 correspond to the death effector domain (DED), which is a domain found in proteins that regulate apoptotic signaling pathways. The DED domain is necessary for PEA-15 to block Ras suppression. Although PEA-15 is predominantly expressed in the central nervous system, low levels of PEA-15 are expressed in liver and kidney, and higher levels in muscle. PEA-15 is also referred to as PED, phosphoprotein enriched in diabetes, for its elevated expression in type 2 diabetic patients

#### Swiss-Prot

Q15121

#### Applications

#### Blocking

#### Specificity

This peptide can be used with studies using BS1573 PEA-15 (D110) pAb.

#### Purification & Purity

Synthetic peptide PEA-15 (D110). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.