

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



PKA α / β cat (V191) Peptide

Cat No.: BS1576P

Background

The second messenger cyclic AMP (cAMP) mediates diverse cellular responses to external signals such as proliferation, ion transport, regulation of metabolism and gene transcription by activation of the cAMP-dependent protein kinase (cAPK or PKA). Activation of PKA occurs when cAMP binds to the two regulatory subunits of the tetrameric PKA holoenzyme resulting in release of active catalytic subunits. Three catalytic (C) subunits have been identified, designated C α , C β and C γ , that each represent specific gene products. C α and C β are closely related (93% amino acid sequence similarity), whereas C γ displays 83% and 79% similarity to C α and C β , respectively.

Swiss-Prot

P17612/P22694

Applications

Blocking

Specificity

This peptide can be used with studies using BS1576 PKA α / β cat (V191) pAb.

Purification & Purity

Synthetic peptide PKA α / β cat (V191). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.