

# PKD1 (R201) polyclonal antibody

Catalog: BS1583

Host: Rabbit

Reactivity: Human, Mouse, Rat

## **BackGround:**

Activation of PKC is one of the earliest events in a cascade leading to a variety of cellular responses such as secretion, gene expression, proliferation and muscle contraction. Protein kinase D (PKD), also called PKCµ, is a serine/threonine kinase whose activation is dependent on the phosphorylation of two activation loop sites, Ser744 and Ser748, via a PKC-dependent signaling pathway. In addition to the two activation loop sites, the carboxy-terminal Ser916 has been identified as an autophosphorylation site for PKD/PKCµ. Phosphorylation at Ser916 correlates with PKD/PKCµ catalytic activity.

**Product:** 

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:** 

~ 115 kDa

**Swiss-Prot:** 

Q15139

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:** 

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at -20  $^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

#### Specificity:

PKD1 (R201) polyclonal antibody detects endogenous levels of PKD1 protein.

**DATA:** 



Immunohistochemistry (IHC) analyzes of PKD1 (R201) pAb in paraffin-embedded human breast tissue.

#### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

## Bioworld technology, co. Ltd.

 
 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151