## **Bioworld Technology CO., Ltd.**



# TH (P2) Peptide

Cat No.: BS1600P

### Background

The enzyme tyrosine hydroxylase (TH), also designated tyrosine 3-monooxygenase (TY3H), catalyzes the conversion of tyrosine to L-dopa, which is the rate limiting step in the biosynthesis of catecholamines such as dopamine, adrenalin and noradrenalin. TH is thought to play a role in the pathogenesis of Parkinson's disease, which is associated with reduced dopamine levels. Two transcription factor binding sites in the proximal region of the TH gene, the TPA-responsive element (TRE) and the c-AMP responsive element (CRE), have been implicated in the complex regulation of the TH gene. TH is also known to be upregulated by the glia maturation factor (GMF), a Cdc 10/SWI6 motif-containing protein called V-1, and a variety of additional compounds

#### **Swiss-Prot**

P07101

Applications

Blocking

#### Specificity

This peptide can be used with studies using BS1600 TH (P2) pAb.

#### **Purification & Purity**

Synthetic peptide TH (P2). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

**Storage & Stability** 

Store at  $4 \, \mathbb{C}$  short term. Aliquot and store at  $-20 \, \mathbb{C}$  long term. Avoid freeze-thaw cycles.

#### **Research Use**

For research use only, not for use in diagnostic procedure.