

PRODUCT DATA SHEET

Bioworld Technology,Inc.

GAK (Q129) polyclonal antibody

Catalog: BS1871 Host: Rabbit Reactivity: Human, Mouse

BackGround:

Cyclin D1 accumulates during G1 and associates with Cdk2, Cdk4 and Cdk5. Cyclin E and Cdk2 interact during the G1 to S transition. Cyclin G contains a typical N terminal cyclin box and a carboxy terminal domain sequence homologous to the tyrosine phosphorylation site of the epidermal growth factor receptor). Cyclin G expression is induced within 3 hours after growth stimulation and remains elevated with no apparent cell cycle dependency. A serine/threonine kinase, designated GAK for cyclin G associated kinase, has been identified. GAK has been shown to bind directly to cyclin G and to co-immunoprecipitate with Cdk5, which also associates with cyclin G.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 143 kDa

Swiss-Prot:

O14976

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:


IHC: 1:50~1:200

Storage&Stability:

Specificity:

GAK (Q129) polyclonal antibody detects endogenous levels of GAK protein.

DATA:



BS1871 Lot CA3613

Immunohistochemistry (IHC) analyzes of GAK (Q129) pAb in paraffin-embedded human kidney carcinoma tissue at 1:50.showing Cytoplasm staining. Negative control (the right)Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151