

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### HMGB2 (K165) Peptide

Cat No.: BS1967P

#### Background

High mobility group (HMG) proteins 1 and 2 are ubiquitous non-histone components of chromatin. Evidence suggests that the binding of HMG proteins to DNA induces alterations in the DNA architecture including DNA bending and unwinding of the helix. HMG proteins synergize with Oct-2, members of the N  $\kappa$ B family, ATF-2 and c-Jun to activate transcription. Other studies indicate that phosphorylation of HMG protein is required to stimulate the transcriptional activity of the protein. Human HMG-1 and HMG-2 both contain two DNA-binding domains, termed HMG boxes. HMG proteins bind single-stranded DNA but induce conformational changes in double-stranded DNA alone.

#### Swiss-Prot

P26583

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS1967 HMGB2 (K165) pAb.

#### Purification & Purity

Synthetic peptide HMGB2 (K165). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.