Bioworld Technology CO., Ltd.



HNF4α/γ (K127) Peptide

Cat No.: BS1992P

Background

HNF-1 (α and β), HNF-3 (α , β and γ), HNF-4 (α and γ), and HNF-6 compose, in part, a homoeprotein family designated the Hepatocyte Nuclear Factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. HNF-3 α , HNF-3 β and HNF-3 γ regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3 α and HNF-3 β have also been shown to be involved in gastrulation events such as body axis formation. HNF-4 α and HNF-4 γ have been shown to be important for early embryo development. HNF-4 α is expressed in liver, kidney, pancreas, small intestine, testis and colon; and HNF-4 γ is expressed in each of these tissues except liver.

Swiss-Prot P41235

Applications

Blocking

Specificity

This peptide can be used with studies using BS1992 HNF4 α/γ (K127) pAb.

Purification & Purity

Synthetic peptide HNF4 α/γ (K127). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.