# **Bioworld Technology CO., Ltd.**



# Cdc20 (Q105) Peptide

Cat No.: BS2203P

# Background

Transition from G1 to S phase requires the association of Cdc28 with members of the G1 cyclin family. Exit from mitosis and initiation of the next cell cycle requires a complex of proteins designated the anaphase-promoting complex (APC). This complex consists of two proteins,Cdc16 and Cdc27 (also referred to as Snb1), which are involved in limiting DNA replication to once per cell cycle. Cdc23, another component of the APC, is required for both entering and exiting anaphase, and is important for the proper separation of sister chromatids. The APC is thought to be stabilized by Cdc26 (also known as Scd26). In addition to these APC proteins, Cdc2 acts as a DNA-damage induced checkpoint, preventing mitosis when DNA damage has occurred..

## **Swiss-Prot**

Q12834

Applications

Blocking

## Specificity

This peptide can be used with studies using BS2203 Cdc20 (Q105) pAb.

#### **Purification & Purity**

Synthetic peptide Cdc20 (Q105). (Note: the amino acid sequence is proprietary). The purity is > 98%.

### Product

1 mg/ml in DI water.

**Storage & Stability** 

Store at  $4 \, \mathbb{C}$  short term. Aliquot and store at  $-20 \, \mathbb{C}$  long term. Avoid freeze-thaw cycles.

#### **Research Use**

For research use only, not for use in diagnostic procedure.