



MEK-4/ MAP2K4 (K350) Peptide

Cat No.: BS2276P

Background

The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK 5, whereas MEK-6 phosphorylates p38 and p38β. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway.

Swiss-Prot

P45985

Applications

Blocking

Specificity

This peptide can be used with studies using BS2276 MEK-4/ MAP2K4 (K350) pAb.

Purification & Purity

Synthetic peptide MEK-4/ MAP2K4 (K350). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.